



CITRUS NOTES

In This Issue:

- Citrus Growers' Educational Seminar
- Management of rind breakdown of mandarins
- Citrus blast

Citrus Growers' Educational Seminar

Sponsored by: University of California Cooperative Extension and Citrus Research Board

When: Wednesday, October 29, 2008

Where: Glenn County Farm Bureau Office - 831 5th Street, Orland

Cost: \$15.00 per person includes lunch and course materials.

Please return attached Advance Registration Form

8:30 Registration Opens

9:00 Introductions & Seminar Sessions Begin

Bill Krueger, UC Farm Advisor and County Director, Glenn County

9:10 Irrigation Scheduling with Microsprinklers: Getting the Most Out of Available Water in a Drought

Larry Schwankl, UC Irrigation Specialist, Kearney Ag. Center

9:40 Principles of Frost Protection

Joe Connell, UC Farm Advisor, Butte County

10:20 Break

10:40 Controlling Brown Rot and Citrus Blast

Bill Krueger, UC Farm Advisor, Glenn County

11:20 Organic Citrus Production

Cindy Fake, UC Farm Advisor, Placer County

12:00 Catered Lunch – pre-registration required

12:45 Reports on Critical Issues: Including the Latest on the Asian Citrus Psyllid and the HLB Threat

Citrus Research Board

1:30 pm Adjourn

**Continuing
Education
Credit
has been
requested**

2279 Del Oro Ave. # B, Oroville, CA 95965 (530) 538-7201 FAX (530) 538-7140 Email: cebutte@ucdavis.edu Web Page: cebutte.ucdavis.edu

University of California, and the United States Department of Agriculture, Cooperating with Butte County

To simplify information trade names of products have been used. No endorsement of named products is intended nor is criticism implied of similar products which are not mentioned...

Inquiries regarding the University's nondiscrimination policies may be directed to the Affirmative Action/Staff Personnel Services Director, University of California, Agriculture and Natural Resources, 1111 Franklin, 6th Floor, Oakland, CA 94607-5200, or phone (510) 987-0096.

Management of rind breakdown of mandarins

By J.E. Adaskaveg

Dept. of Plant Pathology, Univ. of Calif., Riverside

Pre-harvest rind decay of mandarins in California generally occurs shortly after rain falls and is most

severe on Satsuma mandarins. Although some researchers have associated the problem with fungi such as *Alternaria* species, our isolations from affected fruit were inconsistent. Re-inoculations with the isolated fungi (e.g., species of *Alternaria*, *Fusarium*, *Macrophomina*, *Ulocladium*, and *Cladosporium*) only sometimes reproduced disease symptoms and only on water-soaked fruit. Furthermore, in preliminary field trials that were conducted in Butte Co. in the fall of 2002 and 2003, fungicide treatments that included Topsin-M[®], Pristine[®], and Abound[®] only reduced the incidence of disease from 99% in the control to approximately 90%. These data suggested that mandarin rind breakdown is a physiological, abiotic disorder of fruit rather than a pathological problem and the fungi isolated are rather secondary causes of rind decay than primary pathogens.

Rind breakdown of citrus was previously reported by Fawcett and others in the 1930s. Wet weather combined with a sudden decrease in temperature was shown to result in liberation of rind oil and collapse of cells just under the cuticle. In our laboratory studies with Satsuma mandarins, symptoms could be reproduced by water soaking of

the fruit. In laboratory and preliminary field trials in 2003 fruit treatments with water repellants (e.g., postharvest fruit coatings, an agricultural anti-transpirant, or a summer oil) reduced the incidence of rind breakdown to very low levels. Field trials were again conducted in the fall of 2004 in Butte County. Fungicide treatments were ineffective in the Butte Co. trial and in all trials, applications of Vapor-Gard or Omni oil significantly reduced the disorder (Table 1). In all programs with Vapor-Gard and Omni Oil, a first application was made at the end of October and there was no significant difference in efficacy when additional applications were done. When trees were protected from rainfall using a tent, in both 2003 and 2004 (Table 1), the disorder could not be detected indicating the rind breakdown is correlated to rainfall.

In summary, results from our trials support previous findings by Fawcett and others that mandarin rind disorder is an abiotic, weather-related problem of mature fruit that has undergone a green to orange color change. Furthermore, we developed economical treatments that can significantly reduce the disorder.

Table 1. Effect of fungicides and rain protecting materials on mandarin rind disorder in Butte Co. 2004

No.	Treatment*	Product Rates (/200gals/A)	Application Dates				Incidence of MRD (%) & LSD
			Oct. 22	Nov. 9	Nov. 19	Nov. 26	
1	Check	---	---	---	---	---	33.67 A
2	Abound	12.8oz	@	---	@	---	31.25 A
4	Vapor Gard	1gal	@	@	---	@	5.85 B
5	Vapor Gard	1gal	@	---	---	---	4.00 B
7	Omni Oil	6qt-0.75%	@	@	---	@	1.25 BC
8	Omni Oil	6qt-0.75%	@	---	---	---	1.96 BC
9	Vapor Gard/Nordox	1 gal/6 lb	@	---	---	---	5.80 B
10	Tent Protected	---	---	---	---	---	0 C

* - Treatments were applied using an air-blast sprayer calibrated to 200 gal/A.

** - Incidence of mandarin rind disorder (MRD) is based on the evaluation of 100 fruit per tree.

Citrus blast

Blast is caused by the bacterium *Pseudomonas syringae* and is severe only in the northern California counties of Butte, Glenn, and Tehama.

Heavy rains, strong winds, and low temperatures are ideal conditions for citrus blast development. The blast organism usually enters through injured tissues or wounds. Winds that tatter leaves and crack leaf petioles contribute to blast damage.

Thorn, hail or insect injuries to leaves are other avenues of entry for the disease organism.

Blast lesions often start at the leaf petiole. Leaves affected by blast wilt rapidly and dry on the tree. The dried leaves may fall leaving the leaf petiole behind or the entire leaf may remain stuck on the tree. In severe attacks, twigs are girdled and killed. If the twig isn't killed, callus will form at the edge of the twig lesion and the diseased spot will be covered by a reddish brown scab or crust. Blast is most severe on the exposed sides of the tree. In our area, usually the south side and the tree tops are most seriously affected.

The most effective control is a spray before the first rain in late October to early November of a 10-10-100 Bordeaux mixture. Complete coverage using 10-25 gallons of spray per mature tree is the recommended rate. Bordeaux 10-10-100 is made by mixing 10 lbs. copper sulfate (bluestone) containing 25% metallic copper and 10 lbs. fresh

hydrated lime in 100 gallons of water. If you hesitate to spray because Bordeaux is hard to get off the fruit (sometimes a problem with pebbly-skinned mandarins), harvest first and then spray. Commercially available fixed copper materials are not as persistent as a Bordeaux spray. However, they are much easier to use and are readily available.

Sprays applied now will help protect leaves and shoots from winter storms. In years with heavy rainfall, a second spray in late January is appropriate. Remember, a spray application has to be applied before blast injury occurs in order to protect the tissues. Cultural practices that harden off growth in the fall before wet, cold weather comes will help reduce the severity of blast damage. Avoid late season fertilization....the last nitrogen should be applied in late May. Using these cultural and chemical controls will help reduce citrus blast and the severity of blast damage to the tree.

Citrus Growers' Educational Seminar Reservation Form

Advance Reservations are Required

To register, mail coupon no later than October 23rd

or Phone (530) 865-1155, or fax the coupon below to (530) 865-1109

or e-mail to ceglenn@ucdavis.edu by October 23rd. (You may pay at the door.)

----- clip here -----

Please make _____ reservations @ \$15.00 each for the Citrus Growers Educational Seminar to be held on October 29, 2008 in Orland, California:

Name: _____

Address: _____

Phone: _____ Fax: _____ E-Mail: _____

Enclosed is my check in the amount of \$_____. **Make check payable to UC Regents**
Mail check with this form to: Glenn Co. Cooperative Extension, P. O. Box 697, Orland, CA 95963